arXiv:1207.5107v1 [math.NT] 21 Jul 2012

EVALUATION OF THE CONVOLUTION SUMS $\sum_{l+15m=n}\sigma(l)\sigma(m)$ AND $\sum_{3l+5m=n}\sigma(l)\sigma(m)$ AND SOME APPLICATIONS

B. RAMAKRISHNAN AND BRUNDABAN SAHU

Dedicated to Srinivasa Ramanujan on the occasion of his 125th Birth Centenary

Abstract. We evaluate the convolution sums $\sum_{l,m\in\mathbb{N},l+15m=n}\sigma(l)\sigma(m)$ and $\sum_{l,m\in\mathbb{N},3l+5m=n}\sigma(l)\sigma(m)$ for all $n\in\mathbb{N}$ using the theory of quasimodular forms and use these convolution sums to determine the number of representations of a positive integer n by the form

$$x_1^2+x_1x_2+x_2^2+x_3^2+x_3x_4+x_4^2+5(x_5^2+x_5x_6+x_6^2+x_7^2+x_7x_8+x_8^2).$$

We also determine the number of representations of positive integers by the quadratic form

$$x_1^2+x_2^2+x_3^2+x_4^2+6(x_5^2+x_6^2+x_7^2+x_8^2),$$

by using the convolution sums obtained earlier by Alaca, Alaca and Williams [3, 4].

1. Introduction

Following [31, 28], for $n,N\in\mathbb{N}$, we define $W_N(n)$ as follows.

$$W_N(n)=\sum_{m<n/N}\sigma(m)\sigma(n-Nm), \tag{1}$$

where $\sigma_r(n)$ is the sum of the r-th powers of the divisors of n. We write $\sigma_1(n)=\sigma(n)$. Also, following [1], we define $W_{a,b}(n)$ for $a,b\in\mathbb{N}$ by

$$W_{a,b}(n):=\sum_{\substack{l,m\\ al+bm=n}}\sigma(l)\sigma(m). \tag{2}$$

Note that $W_{1,N}(n)=W_{N,1}(n)=W_N(n)$. These type of sums were evaluated as early as the 19th century. For example, the sum $W_1(n)$ was evaluated by Besge, Glaisher and Ramanujan [9, 15, 27].

The convolution sums $W_N(n)$ (for $1\le N\le 24$ with a few exceptions) and $W_{a,b}(n)$ for $(a,b)\in\{(2,3),\ (3,4),(3,8),(2,9)\}$ have been evaluated by using either elementary methods or analytic methods (which use ideas of Ramanujan) or algebraic methods (using quasimodular forms) (cf. [9, 15, 27, 24, 20, 21, 26, 18, 1, 2, 3, 4, 5, 6, 12, 13, 22, 31, 32, 28]). Evaluation of these

---

Date: July 24, 2012.

2000 *Mathematics Subject Classification.* Primary 11A25, 11F11; Secondary 11E20, 11E25, 11F20.

convolution sums have been applied to find the number of representations of integers by certain quadratic forms (cf. [18, 1, 2, 3, 4, 5, 31, 32]). In this article, we evaluate the convolution sums $W_{15}(n)$ and $W_{3,5}(n)$ using the theory of quasimodular forms (following Royer). These convolution sums have not been evaluated so far in the literature (see for example [28, Table 1]). As an application, we use these convolution sums to get the number of representations of integers by the quadratic form $x_1^2+x_1x_2+x_2^2+x_3^2+x_3x_4+x_4^2+5(x_5^2+x_5x_6+x_6^2+x_7^2+x_7x_8+x_8^2)$. We also give a formula for the number of representations of integers by the quadratic forms $x_1^2+x_2^2+x_3^2+x_4^2+k(x_5^2+x_6^2+x_7^2+x_8^2)$, $k=3,6$, by using the convolution sums $W_3(n)$, $W_6(n)$, $W_{12}(n)$, $W_{24}(n)$, $W_{2,3}(n)$ and $W_{3,4}(n)$ evaluated by K. Williams and his co-authors [1, 3, 4, 18].

2. EVALUATION OF $W_{a,b}(n)$ AND SOME APPLICATIONS

2.1. **Evaluation of $W_{15}(n)$ and $W_{3,5}(n)$.**

In this section, following Royer [28], we evaluate the convolution sums $W_{15}(n)$ and $W_{3,5}(n)$ by using the theory of quasimodular forms. As an application, we use these convolution sums together with the convolution sum $W_5(n)$ derived by Lemire and Willams [22] to obtain a formula for the number of representations of a positive integer n by the quadratic form Q given by:

$$Q: x_1^2+x_1x_2+x_2^2+x_3^2+x_3x_4+x_4^2+5(x_5^2+x_5x_6+x_6^2+x_7^2+x_7x_8+x_8^2). \quad (3)$$

Let

$$\Delta_{4,5}(z) = [\Delta(z)\Delta(5z)]^{1/6} = \eta^4(z)\eta^4(5z) = \sum_{n\geq 1} \tau_{4,5}(n)q^n \quad (4)$$

be the normalized newform of weight 4 on $\Gamma_0(5)$ (see [28]). The following theorem was proved by Lemire and Williams [22]:

Theorem 2.1.

$$W_5(n) = \frac{5}{312}\sigma_3(n) + \frac{125}{132}\sigma_3\left(\frac{n}{5}\right) + \frac{5-6n}{120}\sigma(n) + \frac{1-6n}{24}\sigma\left(\frac{n}{5}\right) - \frac{1}{130}\tau_{4,5}(n). \quad (5)$$

In order to evaluate $W_{15}(n)$ and $W_{3,5}(n)$, we use the structure theorem on quasimodular forms of weight k and depth $\leq k/2$. Denoting the space of quasimodular forms of weight k, depth $\leq k/2$ and level N by $\tilde{M}_k^{\leq k/2}\Gamma_0(N))$, we have the following structure theorem (see [19, 25]). For an even integer k with $k\geq 2$, we have

$$\tilde{M}_k^{\leq k/2}(\Gamma_0(N)) = \bigoplus_{i=0}^{k/2-1} D^i M_{k-2i}(\Gamma_0(N)) \oplus \mathbb{C}D^{k/2-1}E_2. \quad (6)$$

Let $\Delta_{4,15;j}(z)$, $j = 1, 2$ be the normalized newforms of weight 4 level 15, which are given by the following.

$$\begin{aligned}\Delta_{4,15;1}(z) &:= \Delta_{4,5}(z) + 9\Delta_{4,5}(3z) + 5\Delta_{2,15}(z)^2 + 2\frac{\eta(z)^5\eta(15z)^5}{\eta(3z)\eta(5z)} \\ &= \sum_{n\geq 1}\tau_{4,15;1}(n)q^n, \\ \Delta_{4,15;2}(z) &:= \Delta_{4,5}(z) + 9\Delta_{4,5}(3z) + 7\Delta_{2,15}(z)^2 = \sum_{n\geq 1}\tau_{4,15;2}(n)q^n,\end{aligned} \tag{7}$$

where

$$\Delta_{2,15}(z) = \eta(z)\eta(3z)\eta(5z)\eta(15z)$$

is the normalized newform of weight 2 and level 15. Using the results of [14, 16] it follows that the functions which are eta-quotients are cusp forms of weight 2 or 4 on $\Gamma_0(15)$. By comparing the first few Fourier coefficients and using the Sturm bound we conclude that they are newforms. The first few Fourier coefficients of newforms of given weight and level are obtained using the database of L-functions, modular forms, and related objects (see [23]).

The following are the main theorems of this section.

Theorem 2.2. *Let $n \in \mathbb{N}$, then*

$$\begin{aligned}W_{15}(n) \;=\;& \frac{5-2n}{120}\sigma(n) + \frac{1-6n}{24}\sigma\left(\frac{n}{15}\right) + \frac{1}{624}\sigma_3(n) + \frac{3}{208}\sigma_3\left(\frac{n}{3}\right) \\ &+\frac{25}{624}\sigma_3\left(\frac{n}{5}\right) + \frac{75}{208}\sigma_3\left(\frac{n}{15}\right) - \frac{1}{455}\tau_{4,5}(n) - \frac{9}{455}\tau_{4,5}\left(\frac{n}{3}\right) \\ &-\frac{1}{84}\tau_{4,15;1}(n) - \frac{1}{80}\tau_{4,15;2}(n), \\ W_{3,5}(n) \;=\;& \frac{5-6n}{120}\sigma(\frac{n}{3}) + \frac{1-2n}{24}\sigma\left(\frac{n}{5}\right) + \frac{1}{624}\sigma_3(n) + \frac{3}{208}\sigma_3\left(\frac{n}{3}\right) \\ &+\frac{25}{624}\sigma_3\left(\frac{n}{5}\right) + \frac{75}{208}\sigma_3\left(\frac{n}{15}\right) - \frac{1}{455}\tau_{4,5}(n) - \frac{9}{455}\tau_{4,5}\left(\frac{n}{3}\right) \\ &-\frac{1}{84}\tau_{4,15;1}(n) + \frac{1}{80}\tau_{4,15;2}(n).\end{aligned}$$

Proof. Using the structure of $\tilde{M}_4^{\leq 2}(15)$ from (6), we get

$$\tilde{M}_4^{\leq 2}(\Gamma_0(15)) = M_4(\Gamma_0(15)) \oplus DM_2(\Gamma_0(15)) \oplus \mathbb{C}DE_2. \tag{8}$$

The vector space $M_4(\Gamma_0(15))$ has dimension 8 and is spanned by the eight linearly independent modular forms $E_4(z), E_4(3z), E_4(5z), E_4(15z), \Delta_{4,5}(z)$, $\Delta_{4,5}(3z)$ and the two normalized newforms $\Delta_{4,15;1}$ and $\Delta_{4,15;2}$. The vector space $M_2(\Gamma_0(15))$ has dimension 4 and is spanned by the linearly independent modular forms $\Phi_{1,15}(z), \Phi_{5,15}(z), \Phi_{1,3}(z)$ and the normalized newform $\Delta_{2,15}(z)$, where

$$\Phi_{a,b}(z) := \frac{1}{b-a}(bE_2(bz) - aE_2(az)). \tag{9}$$

Consider the quasimodular form $E_2(z)E_2(15z)$ which belongs to $\tilde{M}_4^{\leq 2}(\Gamma_0(15))$. Therefore, using (6), we have

$$\begin{aligned} E_2(z)E_2(15z) &= \frac{1}{260}E_4(z) + \frac{9}{260}E_4(3z) + \frac{5}{52}E_4(5z) + \frac{45}{52}E_4(15z) \\ &\quad -\frac{576}{455}\Delta_{4,5}(z) - \frac{5184}{455}\Delta_{4,5}(3z) - \frac{48}{7}\Delta_{4,15;1}(z) \\ &\quad -\frac{36}{5}\Delta_{4,15;2}(z) + \frac{28}{5}D\Phi_{1,15}(z) + \frac{4}{5}DE_2(z). \end{aligned}$$

Similarly, considering $E_2(3z)E_2(5z)$, which is a quasimodular form of weight 4, depth 2 and level 15, we get

$$\begin{aligned} E_2(3z)E_2(5z) &= \frac{1}{260}E_4(z) + \frac{9}{260}E_4(3z) + \frac{5}{52}E_4(5z) + \frac{45}{52}E_4(15z) \\ &\quad -\frac{576}{455}\Delta_{4,5}(z) - \frac{5184}{455}\Delta_{4,5}(3z) - \frac{48}{7}\Delta_{4,15;1}(z) + \frac{36}{5}\Delta_{4,15;2}(z) \\ &\quad +\frac{28}{5}D\Phi_{1,15}(z) - 4D\Phi_{5,15}(z) + \frac{4}{5}D\Phi_{1,3}(z) + \frac{4}{5}DE_2(z). \end{aligned}$$

By comparing the n-th Fourier coefficients, we get the required the convolution sums. □

2.2. **Application to the number of representations.**

In this section we apply the convolution sums $W_{15}(n)$ and $W_{3,5}(n)$ to derive the following theorem.

Theorem 2.3. *The number of representations of a positive integer n by the quadratic form $x_1^2+x_1x_2+x_2^2+x_3^2+x_3x_4+x_4^2+5(x_5^2+x_5x_6+x_6^2+x_7^2+x_7x_8+x_8^2)$ is equal to*

$$\begin{aligned} \frac{12}{13}\sigma_3(n) + \frac{108}{13}\sigma_3\left(\frac{n}{3}\right) + \frac{300}{13}\sigma_3\left(\frac{n}{5}\right) + \frac{2700}{13}\sigma_3\left(\frac{n}{15}\right) + \frac{72}{91}\tau_{4,5}(n) \\ +\frac{648}{91}\tau_{4,5}\left(\frac{n}{3}\right) + \frac{72}{7}\tau_{4,15;1}(n). \end{aligned}$$

Proof. Let $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For $l \in \mathbb{N}_0$, let

$$r(l) = \#\left\{(x_1,x_2,x_3,x_4) \in \mathbb{Z}^4 | x_1^2 + x_1x_2 + x_2^2 + x_3^2 + x_3x_4 + x_4^2 = l\right\}$$

so that $r(0) = 1$. For $l \in \mathbb{N}$, we know that (see [11, 18])

$$r(l) = 12\sum_{\substack{d|l, \\ 3\nmid d}} d = 12\sigma(l) - 36\sigma\left(\frac{l}{3}\right).$$

Let $N(n)$ be the number of representations of the given quadratic form Q defined by (3). Then $N(n)$ is given by

$$
\begin{aligned}
N(n) &= \sum_{\substack{l.m\in\mathbb{N}_0\\ l+5m=n}} \left(\sum_{\substack{(x_1,x_2,x_3,x_4)\in\mathbb{Z}^4\\ x_1^2+x_1x_2+x_2^2+x_3^2+x_3x_4+x_4^2=l}} 1 \right) \left(\sum_{\substack{(x_5,x_6,x_7,x_8)\in\mathbb{Z}^4\\ x_5^2+x_5x_6+x_6^2+x_7^2+x_7x_8+x_8^2=m}} 1 \right)\\
&= r(0)r\left(\frac{n}{5}\right) + r(n)r(0) + \sum_{\substack{l,m\in\mathbb{N}\\ l+5m=n}} r(l)r(m)\\
&= 12\sigma\left(\frac{n}{5}\right) - 36\sigma\left(\frac{n}{15}\right) + 12\sigma(n) - 36\sigma\left(\frac{n}{3}\right)\\
&\quad + \sum_{\substack{l,m\in\mathbb{N}\\ l+5m=n}} \left(12\sigma(l) - 36\sigma\left(\frac{l}{3}\right)\right)\left(12\sigma(m) - 36\sigma\left(\frac{m}{3}\right)\right)\\
&= 12\sigma\left(\frac{n}{5}\right) - 36\sigma\left(\frac{n}{15}\right) + 12\sigma(n) - 36\sigma\left(\frac{n}{3}\right) + 144 \sum_{\substack{l,m\in\mathbb{N}\\ l+5m=n}} \sigma(l)\sigma(m)\\
&\quad -432 \sum_{\substack{l,m\in\mathbb{N}\\ l+5m=n}} \sigma(l)\sigma\left(\frac{m}{3}\right) - 432 \sum_{\substack{l,m\in\mathbb{N}\\ l+5m=n}} \sigma\left(\frac{l}{3}\right)\sigma(m)\\
&\quad +1296 \sum_{\substack{l,m\in\mathbb{N}\\ l+5m=n}} \sigma\left(\frac{l}{3}\right)\sigma\left(\frac{m}{3}\right)\\
&= 12\sigma\left(\frac{n}{5}\right) - 36\sigma\left(\frac{n}{15}\right) + 12\sigma(n) - 36\sigma\left(\frac{n}{3}\right)\\
&\quad +144\, W_5(n) - 432\, W_{15}(n) - 432\, W_{3,5}(n) + 1296\, W_5\left(\frac{n}{3}\right).
\end{aligned}
$$

Substituting the convolution sums using Theorem 2.1 and Theorem 2.2, we get the required formula for $N(n)$. $\square$

2.3. **More applications.** Let Q_k be the quadratic form $x_1^2+x_2^2+x_3^2+x_4^2+k(x_5^2+x_6^2+x_7^2+x_8^2)$ and $N_k(n)$ be the number of representations of integers $n \geq 1$ by Q_k. In this section we use the convolution sums derived in [3, 4] to derive a formula for N_6. We note that for $k = 2, 3, 4$ similar formulas were obtained earlier by Williams [32], Alaca-Williams [7] and Alaca-Alaca-Williams [5] respectively. To find $N_6(n)$, we need the convolution sums $W_6(n)$, $W_{2,3}$ and $W_{24}(n)$ which were derived by Alaca-Alaca-Williams and they are given in the following theorem.

Theorem 2.4. (cf. [3, 4])

$$\begin{aligned}
W_6(n) &= \frac{1}{120}\sigma_3(n)+\frac{1}{30}\sigma_3\left(\frac{n}{2}\right)+\frac{3}{40}\sigma_3\left(\frac{n}{3}\right)+\frac{3}{10}\sigma_3\left(\frac{n}{6}\right)\\
&\quad+\frac{1-n}{24}\sigma(n)+\frac{1-6n}{24}\sigma\left(\frac{n}{6}\right)-\frac{1}{120}c_6(n),\\
W_{2,3}(n) &= \frac{1}{120}\sigma_3(n)+\frac{1}{30}\sigma_3\left(\frac{n}{2}\right)+\frac{3}{40}\sigma_3\left(\frac{n}{3}\right)+\frac{3}{10}\sigma_3\left(\frac{n}{6}\right)\\
&\quad+\frac{1-2n}{24}\sigma\left(\frac{n}{2}\right)+\frac{1-3n}{24}\sigma\left(\frac{n}{3}\right)-\frac{1}{120}c_6(n),\\
W_{24}(n) &= \frac{1}{1920}\sigma_3(n)+\frac{1}{640}\sigma_3\left(\frac{n}{2}\right)+\frac{3}{640}\sigma_3\left(\frac{n}{3}\right)+\frac{1}{160}\sigma_3\left(\frac{n}{4}\right)\\
&\quad+\frac{9}{640}\sigma_3\left(\frac{n}{6}\right)+\frac{1}{30}\sigma_3\left(\frac{n}{8}\right)+\frac{9}{160}\sigma_3\left(\frac{n}{12}\right)+\frac{3}{10}\sigma_3\left(\frac{n}{24}\right)\\
&\quad+\frac{4-n}{96}\sigma(n)+\frac{1-6n}{24}\sigma\left(\frac{n}{24}\right)-\frac{61}{1920}c_{1,24}(n),
\end{aligned}$$

where $c_6(n)$ *and* $c_{1,24}(n)$ *are the* n*-th Fourier coefficients of weight* 4 *normalized newforms which are given in* [3, p. 492] *and* [4, p. 94] *respectively.*

In the following we use Theorem 2.4 to derive a formula for $N_6(n)$.

Theorem 2.5. *The number of representations of a positive integer* n *by the quadratic form* Q_6 *is given by*

$$\begin{aligned}
N_6(n) &= \frac{2}{5}\sigma_3(n)-\frac{2}{5}\sigma_3\left(\frac{n}{2}\right)+\frac{18}{5}\sigma_3\left(\frac{n}{3}\right)-\frac{8}{5}\sigma_3\left(\frac{n}{4}\right)-\frac{18}{5}\sigma_3\left(\frac{n}{6}\right)\\
&\quad+\frac{128}{5}\sigma_3\left(\frac{n}{8}\right)-\frac{72}{5}\sigma_3\left(\frac{n}{12}\right)+\frac{1152}{5}\sigma_3\left(\frac{n}{24}\right)-\frac{8}{15}c_6(n)\\
&\quad+\frac{32}{15}c_6\left(\frac{n}{2}\right)-\frac{128}{15}c_6\left(\frac{n}{4}\right)+\frac{122}{15}c_{1,24}(n).
\end{aligned}$$

Proof. For $l \in \mathbb{N}_0$, let

$$r_4(l) = \#\left\{(x_1,x_2,x_3,x_4)\in\mathbb{Z}^4 | x_1^2+x_2^2+x_3^2+x_4^2=l\right\}$$

so that $r(0)=1$. For $l\in\mathbb{N}$, we know the formula due to Jacobi (see [17])

$$r_4(l) = 8\sum_{\substack{d|l,\\ 4\nmid d}} d = 8\sigma(l)-32\sigma\left(\frac{l}{4}\right).$$

Then N_6 is given by

$$\begin{aligned}
N_6(n) &= \sum_{\substack{l.m\in\mathbb{N}_0\\ l+6m=n}} \left(\sum_{\substack{(x_1,x_2,x_3,x_4)\in\mathbb{Z}^4\\ x_1^2+x_2^2+x_3^2+x_4^2=l}} 1 \right) \left(\sum_{\substack{(x_5,x_6,x_7,x_8)\in\mathbb{Z}^4\\ x_5^2+x_6^2+x_7^2+x_8^2=m}} 1 \right) \\
&= r_4(0)r_4\left(\frac{n}{6}\right) + r_4(n)r_4(0) + \sum_{\substack{l,m\in\mathbb{N}\\ l+6m=n}} r_4(l)r_4(m) \\
&= 8\sigma\,(n) - 32\sigma\left(\frac{n}{4}\right) + 8\sigma\left(\frac{n}{6}\right) - 32\sigma\left(\frac{n}{24}\right) \\
&\quad + \sum_{\substack{l,m\in\mathbb{N}\\ l+6m=n}} \left(8\sigma(l) - 32\sigma\left(\frac{l}{4}\right)\right)\left(8\sigma(m) - 32\sigma\left(\frac{m}{4}\right)\right) \\
&= 8\sigma\,(n) - 32\sigma\left(\frac{n}{4}\right) + 8\sigma\left(\frac{n}{6}\right) - 32\sigma\left(\frac{n}{24}\right) + 64\sum_{\substack{l,m\in\mathbb{N}\\ l+6m=n}} \sigma(l)\sigma(m) \\
&\quad -256\sum_{\substack{l,m\in\mathbb{N}\\ l+6m=n}} \sigma(l)\sigma\left(\frac{m}{4}\right) - 256\sum_{\substack{l,m\in\mathbb{N}\\ l+6m=n}} \sigma\left(\frac{l}{4}\right)\sigma(m) \\
&\quad +1024\sum_{\substack{l,m\in\mathbb{N}\\ l+6m=n}} \sigma\left(\frac{l}{4}\right)\sigma\left(\frac{m}{4}\right) \\
&= 8\sigma\,(n) - 32\sigma\left(\frac{n}{4}\right) + 8\sigma\left(\frac{n}{6}\right) - 32\sigma\left(\frac{n}{24}\right) + 64\;W_6(n) \\
&\quad +1024\;W_6\left(\frac{n}{4}\right) - 256\;W_{24}(n) - 256\;W_{2,3}\left(\frac{n}{2}\right).
\end{aligned}$$

Substituting the convolution sums from Theorem 2.4 in the above gives the required formula for $N_6(n)$. $\square$

Remark 2.1. The representation numbers $N_k(n)$ for $k = 2, 4$ were obtained by Williams [32] and by Alaca-Alaca-Williams [5] using the convolution sums $W_{a,b}(n)$ and for $k = 3$ it was derived by Alaca-Williams [7] as a consequence of the representation of positive integers by certain octonary quadratic forms. Note that $N_3(n)$ can also be obtained in a similar way as done in the cases $k = 2, 4$. In fact,

$$\begin{aligned}
N_3(n) &= 8\sigma(n) - 32\sigma\left(\frac{n}{4}\right) + 8\sigma\left(\frac{n}{3}\right) - 32\sigma\left(\frac{n}{12}\right) \qquad (10) \\
&\quad +64\;W_3(n) + 1024\;W_3\left(\frac{n}{4}\right) - 256\;W_{12}(n) - 256\;W_{3,4}(n).
\end{aligned}$$

Using the convolution sums $W_3(n)$, $W_{3,4}(n)$ and $W_{12}(n)$ obtained in [18, 1], we have the following formula for $N_3(n)$:

$$\begin{aligned} N_3(n) \;\; = \;\; & \frac{8}{5}\,\sigma_3(n) - \frac{16}{5}\,\sigma_3\left(\frac{n}{2}\right) + \frac{72}{5}\,\sigma_3\left(\frac{n}{3}\right) + \frac{128}{5}\,\sigma_3\left(\frac{n}{4}\right) \\ & -\frac{144}{5}\,\sigma_3\left(\frac{n}{6}\right) + \frac{1152}{5}\,\sigma_3\left(\frac{n}{12}\right) + \frac{88}{15}\,c_{1,12}(n) + \frac{8}{15}\,c_{3,4}(n). \end{aligned} \tag{11}$$

The difference between the formula given in [7, Theorem 1.1 (ii)] and (11) is the coefficients coming from cusp forms. In [7] coefficients of the newform of weight 4 level 6 appears and in the above formula Fourier coefficients of two cusp forms of weight 4 and level 12 appear.

Acknowledgements. We have used the open-source mathematics software SAGE (www.sagemath.org) to do our calculations. The work was done during the second author's visit to the Harish-Chandra Research Institute (HRI), Allahabad. He wishes to thank HRI for the warm hospitality during his stay.

References

[1] A. Alaca, S. Alaca and K. S. Williams, *Evaluation of the convolution sums* $\sum_{l+12m=n}\sigma(l)\sigma(m)$ *and* $\sum_{3l+4m=n}\sigma(l)\sigma(m)$, Adv. Theor. Appl. Math. **1** (2006), no. 1, 27–48.

[2] A. Alaca, S. Alaca and K. S. Williams,*Evaluation of the convolution sums* $\sum_{l+18m=n}\sigma(l)\sigma(m)$ *and* $\sum_{2l+9m=n}\sigma(l)\sigma(m)$, Int. Math. Forum **2** (2007), no. 1-4, 45–68.

[3] A. Alaca, S. Alaca and K. S. Williams, *Evaluation of the convolution sums* $\sum_{l+6m=n}\sigma(l)\sigma(m)$ *and* $\sum_{2l+3m=n}\sigma(l)\sigma(m)$, J. Number Theory **124** (2007), no. 2, 491–510.

[4] A. Alaca, S. Alaca and K. S. Williams, *Evaluation of the convolution sums* $\sum_{l+24m=n}\sigma(l)\sigma(m)$ *and* $\sum_{3l+8m=n}\sigma(l)\sigma(m)$, Math. J. Okayama Univ. **49** (2007), 93–111.

[5] A. Alaca, S. Alaca and K. S. Williams, *The convolution sum* $\sum_{m<n/16}\sigma(m)\sigma(n-16m)$, Canad. Math. Bull. **51** (2008), no. 1, 3–14.

[6] A. Alaca, S. Alaca, F. Uygul and K. S. Williams, *Restricted Eisenstein series and certain convolution sums*, J. Comb. Number Theory **3** (2011), 1–14.

[7] S. Alaca, K. Williams, *The number of representations of a positive integer by certain octonary quadratic forms*, Funct. Approx. Comment. Math. **43** (2010), part 1, 45-54.

[8] A. O. L. Atkin and J. Lehner, *Hecke operators on* $\Gamma_0(m)$, Math. Ann. **185** (1970), 134–160.

[9] M. Besge, *Extrait d'une lettre de M. Besge á M. Liouville,* J. Math. Pures Appl. **7** (1862), 256.

[10] J. H. Bruinier, G. van der Geer, G. Harder and D. Zagier, *The 1-2-3 of modular forms*, Lectures from the Summer School on Modular Forms and their Applications held in Nordfjordeid, June 2004. Edited by Kristian Ranestad. Universitext. Springer-Verlag, Berlin, 2008. 266 pp.

[11] R. Chapman, *Representations of integers by the form* $x^2+xy+y^2+z^2+zt+t^2$, Int. J. Number Theory **4** (2008), no. 5, 709–714.

[12] N. Cheng and K. S. Williams, *Convolution sums involving the divisor function*, Proc. Edinb. Math. Soc. (2) **47** (2004), no. 3, 561–572.

[13] N. Cheng and K. S. Williams, *Evaluation of some convolution sums involving the sum of divisors functions*, Yokohama Math. J. **52** (2005), no. 1, 39–57.

[14] D. Dummit, H. Kisilesky and J. McKay, Multiplicative products of η-functions. Finite groupscoming of age (Montreal, Que., 1982), 8998, Contemp. Math., 45, Amer. Math. Soc., Providence, RI, 1985.

[15] J. W. L. Glaisher, *On the squares of the series in which the coefficients are the sums of the divisor of the exponents*, Mess. Math. **15** (1885), 1–20.

[16] B. Gordon and D. Sinor, Multiplicative properties of η-products. Number theory, Madras 1987, 173200, Lecture Notes in Math., 1395, Springer, Berlin, 1989.

[17] E. Grosswald, Representation of integers as sums of squares, *Springer, New York*, 1985

[18] J. G. Huard, Z. M. Ou, B. K. Spearman and K. S. Williams, *Elementary evaluation of certain convolution sums involving divisor functions,* in Number Theory for the Millennium, II (Urbana, IL, 2000) (A. K. Peters, Natick, MA, 2002), 229–274.

[19] M. Kaneko and D. Zagier, *A generalized Jacobi theta function and quasimodular forms.* In 'The moduli space of curves (Texel Island, 1994)', 165–172, Progr. Math., **129**, Birkhäuser Boston, Boston, MA, 1995.

[20] D. B. Lahiri, *On Ramanujan's function $\tau(n)$ and the divisor function $\sigma_k(n)$ - I*, Bull. Calcutta Math. Soc. **38** (1946), 193–206.

[21] D. B. Lahiri, *On Ramanujan's function $\tau(n)$ and the divisor function $\sigma_k(n)$ - II*, Bull. Calcutta Math. Soc. **39** (1947), 33–52.

[22] M. Lemire and K. S. Williams, *Evaluation of two convolution sums involving the sum of divisors function*, Bull. Austral. Math. Soc. **73** (2006), no. 1, 107–115.

[23] LMFDB, The database of L-functions, modular forms, and related objects, http://www.lmfdb.org/

[24] P. A. MacMahon, *Divisors of numbers and their continuations in the theory of partitions*, Proc. London Math. Soc., (2) **19** (1920), 75–113.

[25] F. Martin and E. Royer, *Formes modulaires et périodes*, In 'Formes modulaires et transcendance', 1–117, Sémin. Congr., **12**, Soc. Math. France, Paris, 2005.

[26] G. Melfi, *On some modular identities*, in Number Theory (Eger, 1996) (de Gruyter, Berlin, 1998), pp. 371–382;

[27] S. Ramanujan, *On certain arithmetical functions,* Trans. Cambridge Philos. Soc. 22 (1916) 159–184.

[28] E. Royer, *Evaluating convolution sums of the divisor function by quasimodular forms*, Int. J. Number Theory **3** (2007), no. 2, 231–261.

[29] J.-P. Serre, A course in Arithmetic, Springer-Verlag, Berlin-New York, 1977.

[30] G. Shimura, Introduction to the arithmetic theory of automorphic functions. (English summary) Reprint of the 1971 original. Publications of the Mathematical Society of Japan, 11. Kanô Memorial Lectures, 1. Princeton University Press, Princeton, NJ, 1994.

[31] K. S. Williams, *The convolution sum* $\sum_{m<n/9}\sigma(m)\sigma(n-9m)$, Int. J. Number Theory **1** (2005), no. 2, 193–205.

[32] K. S. Williams, *The convolution sum* $\sum_{m<n/8}\sigma(m)\sigma(n-8m)$, Pacific J. Math. **228** (2006), no. 2, 387–396.

(B. Ramakrishnan) Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Allahabad - 211 019, India.

(Brundaban Sahu) School of Mathematical Sciences, National Institute of Science Education and Research, PO: Sainik School, Bhubaneswar, Odisha - 751 005, India.

E-mail address, B. Ramakrishnan: `ramki@hri.res.in`

E-mail address, Brundaban Sahu: `brundaban.sahu@niser.ac.in`